

Hemorrhoids

National Digestive Diseases Information Clearinghouse



U.S. Department
of Health and
Human Services


NATIONAL
INSTITUTES
OF HEALTH

NIDDK
NATIONAL INSTITUTE OF
DIABETES AND DIGESTIVE
AND KIDNEY DISEASES

What are hemorrhoids?

Hemorrhoids are swollen and inflamed veins around the anus or in the lower rectum. The rectum is the last part of the large intestine leading to the anus. The anus is the opening at the end of the digestive tract where bowel contents leave the body.

External hemorrhoids are located under the skin around the anus. Internal hemorrhoids develop in the lower rectum. Internal hemorrhoids may protrude, or prolapse, through the anus. Most prolapsed hemorrhoids shrink back inside the rectum on their own. Severely prolapsed hemorrhoids may protrude permanently and require treatment.



Hemorrhoids

What are the symptoms of hemorrhoids?

The most common symptom of internal hemorrhoids is bright red blood on stool, on toilet paper, or in the toilet bowl after a bowel movement. Internal hemorrhoids that are not prolapsed are usually not painful. Prolapsed hemorrhoids often cause pain, discomfort, and anal itching.

Blood clots may form in external hemorrhoids. A blood clot in a vein is called a thrombosis. Thrombosed external hemorrhoids cause bleeding, painful swelling, or a hard lump around the anus. When the blood clot dissolves, extra skin is left behind. This skin can become irritated or itch.

Excessive straining, rubbing, or cleaning around the anus may make symptoms, such as itching and irritation, worse.

Hemorrhoids are not dangerous or life threatening. Symptoms usually go away within a few days, and some people with hemorrhoids never have symptoms.

How common are hemorrhoids?

About 75 percent of people will have hemorrhoids at some point in their lives.¹ Hemorrhoids are most common among adults ages 45 to 65.² Hemorrhoids are also common in pregnant women.

What causes hemorrhoids?

Swelling in the anal or rectal veins causes hemorrhoids. Several factors may cause this swelling, including

- chronic constipation or diarrhea
- straining during bowel movements
- sitting on the toilet for long periods of time
- a lack of fiber in the diet

Another cause of hemorrhoids is the weakening of the connective tissue in the rectum and anus that occurs with age.

Pregnancy can cause hemorrhoids by increasing pressure in the abdomen, which may enlarge the veins in the lower rectum and anus. For most women, hemorrhoids caused by pregnancy disappear after childbirth.

How are hemorrhoids diagnosed?

The doctor will examine the anus and rectum to determine whether a person has hemorrhoids. Hemorrhoid symptoms are similar to the symptoms of other anorectal problems, such as fissures, abscesses, warts, and polyps.

The doctor will perform a physical exam to look for visible hemorrhoids. A digital rectal exam with a gloved, lubricated finger and an anoscope—a hollow, lighted tube—may be performed to view the rectum.

A thorough evaluation and proper diagnosis by a doctor is important any time a person notices bleeding from the rectum or blood in the stool. Bleeding may be a symptom of other digestive diseases, including colorectal cancer.

Additional exams may be done to rule out other causes of bleeding, especially in people age 40 or older:

- **Colonoscopy.** A flexible, lighted tube called a colonoscope is inserted through the anus, the rectum, and the upper part of the large intestine, called the colon. The colonoscope transmits images of the inside of the rectum and the entire colon.
- **Sigmoidoscopy.** This procedure is similar to colonoscopy, but it uses a shorter tube called a sigmoidoscope and transmits images of the rectum and the sigmoid colon, the lower portion of the colon that empties into the rectum.
- **Barium enema x ray.** A contrast material called barium is inserted into the colon to make the colon more visible in x-ray pictures.

¹Baker H. Hemorrhoids. In: Longe JL, ed. *Gale Encyclopedia of Medicine*. 3rd ed. Detroit: Gale; 2006: 1766–1769.

²Chong PS, Bartolo DCC. Hemorrhoids and fissure in ano. *Gastroenterology Clinics of North America*. 2008;37:627–644.

How are hemorrhoids treated?

At-home Treatments

Simple diet and lifestyle changes often reduce the swelling of hemorrhoids and relieve hemorrhoid symptoms. Eating a high-fiber diet can make stools softer and easier to pass, reducing the pressure on hemorrhoids caused by straining.

Fiber is a substance found in plants. The human body cannot digest fiber, but fiber helps improve digestion and prevent constipation. Good sources of dietary fiber are fruits, vegetables, and whole grains. On average, Americans eat about 15 grams of fiber each day.³ The American Dietetic Association recommends 25 grams of fiber per day for women and 38 grams of fiber per day for men.³

Doctors may also suggest taking a bulk stool softener or a fiber supplement such as psyllium (Metamucil) or methylcellulose (Citrucel).

Other changes that may help relieve hemorrhoid symptoms include

- drinking six to eight 8-ounce glasses of water or other nonalcoholic fluids each day
- sitting in a tub of warm water for 10 minutes several times a day
- exercising to prevent constipation
- not straining during bowel movements

Over-the-counter creams and suppositories may temporarily relieve the pain and itching of hemorrhoids. These treatments should only be used for a short time because long-term use can damage the skin.

Medical Treatment

If at-home treatments do not relieve symptoms, medical treatments may be needed. Outpatient treatments can be performed in a doctor's office or a hospital. Outpatient treatments for internal hemorrhoids include the following:

- **Rubber band ligation.** The doctor places a special rubber band around the base of the hemorrhoid. The band cuts off circulation, causing the hemorrhoid to shrink. This procedure should be performed only by a doctor.
- **Sclerotherapy.** The doctor injects a chemical solution into the blood vessel to shrink the hemorrhoid.
- **Infrared coagulation.** The doctor uses heat to shrink the hemorrhoid tissue.




Large external hemorrhoids or internal hemorrhoids that do not respond to other treatments can be surgically removed.

³Slavin JL. Position statement of the American Dietetic Association: health implications of dietary fiber. *Journal of the American Dietetic Association*. 2008;108(10):1716-1731.

What foods have fiber?

Examples of foods that have fiber include

Beans, cereals, and breads	Fiber
1/2 cup of navy beans	9.5 grams
1/2 cup of kidney beans	8.2 grams
1/2 cup of black beans	7.5 grams
Whole-grain cereal, cold	
1/2 cup of All-Bran	9.6 grams
3/4 cup of Total	2.4 grams
3/4 cup of Post Bran Flakes	5.3 grams
1 packet of whole-grain cereal, hot (oatmeal, Wheatena)	3.0 grams
1 whole-wheat English muffin	4.4 grams
Fruits	
1 medium apple, with skin	3.3 grams
1 medium pear, with skin	4.3 grams
1/2 cup of raspberries	4.0 grams
1/2 cup of stewed prunes	3.8 grams
Vegetables	
1/2 cup of winter squash	2.9 grams
1 medium sweet potato with skin	4.8 grams
1/2 cup of green peas	4.4 grams
1 medium potato with skin	3.8 grams
1/2 cup of mixed vegetables	4.0 grams
1 cup of cauliflower	2.5 grams
1/2 cup of spinach	3.5 grams
1/2 cup of turnip greens	2.5 grams



Source: U.S. Department of Agriculture and U.S. Department of Health and Human Services, *Dietary Guidelines for Americans*, 2005.

Points to Remember

- Hemorrhoids are swollen and inflamed veins around the anus or in the lower rectum.
- Hemorrhoids are not dangerous or life threatening, and symptoms usually go away within a few days.
- A thorough evaluation and proper diagnosis by a doctor is important any time a person notices bleeding from the rectum or blood in the stool.
- Simple diet and lifestyle changes often reduce the swelling of hemorrhoids and relieve hemorrhoid symptoms.
- If at-home treatments do not relieve symptoms, medical treatments may be needed.

Hope through Research

The National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK) sponsors research to improve treatment for people with digestive disorders, including hemorrhoids and constipation. Researchers are studying new drugs and surgical procedures to treat or prevent hemorrhoids.

Participants in clinical trials can play a more active role in their own health care, gain access to new research treatments before they are widely available, and help others by contributing to medical research. For information about current studies, visit www.ClinicalTrials.gov.

For More Information

American College of Gastroenterology
P.O. Box 342260
Bethesda, MD 20827-2260
Phone: 301-263-9000
Internet: www.acg.gi.org

American Gastroenterological Association
4930 Del Ray Avenue
Bethesda, MD 20814
Phone: 301-654-2055
Fax: 301-654-5920
Email: member@gastro.org
Internet: www.gastro.org

American Society of Colon and Rectal Surgeons
85 West Algonquin Road, Suite 550
Arlington Heights, IL 60005
Phone: 847-290-9184
Fax: 847-290-9203
Email: ascrs@fascrs.org
Internet: www.fascrs.org

Acknowledgments

Publications produced by the Clearinghouse are carefully reviewed by both NIDDK scientists and outside experts. This fact sheet was reviewed by Adil E. Bharucha, M.B.B.S., M.D., Mayo Clinic, Rochester, MN.

Saratoga Schenectady
Endoscopy Center, LLC
Burnt Hills, N.Y. 12027

You may also find additional information about this topic by visiting MedlinePlus at www.medlineplus.gov.

This publication may contain information about medications. When prepared, this publication included the most current information available. For updates or for questions about any medications, contact the U.S. Food and Drug Administration toll-free at 1-888-INFO-FDA (1-888-463-6332) or visit www.fda.gov. Consult your doctor for more information.

The U.S. Government does not endorse or favor any specific commercial product or company. Trade, proprietary, or company names appearing in this document are used only because they are considered necessary in the context of the information provided. If a product is not mentioned, the omission does not mean or imply that the product is unsatisfactory.

National Digestive Diseases Information Clearinghouse

2 Information Way

Bethesda, MD 20892-3570

Phone: 1-800-891-5389

TTY: 1-866-569-1162

Fax: 703-738-4929

Email: nddic@info.niddk.nih.gov

Internet: www.digestive.niddk.nih.gov

The National Digestive Diseases Information Clearinghouse (NDDIC) is a service of the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK). The NIDDK is part of the National Institutes of Health of the U.S. Department of Health and Human Services. Established in 1980, the Clearinghouse provides information about digestive diseases to people with digestive disorders and to their families, health care professionals, and the public. The NDDIC answers inquiries, develops and distributes publications, and works closely with professional and patient organizations and Government agencies to coordinate resources about digestive diseases.

This publication is not copyrighted. The Clearinghouse encourages users of this publication to duplicate and distribute as many copies as desired.

This publication is available at
www.digestive.niddk.nih.gov.



U.S. DEPARTMENT OF HEALTH
AND HUMAN SERVICES
National Institutes of Health

NIH Publication No. 11-3021
November 2010



The NIDDK prints on recycled paper with bio-based ink.